

## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



Reserve  
aSB205  
.S7N5

U.S. Department of Agriculture  
Reference Center  
St., S.W., Rm. 147  
Washington, D.C. 20250

ESCS REFERENCE CENTER  
R  
JUN 8 1982

# STAFF REPORT

1980 PESTICIDE USE ON SOYBEANS  
IN THE SOUTHEAST

by

Michael Hanthorn, Craig Osteen,  
Robert McDowell, and Larry Roberson

February 1982


ERS Staff Report No. AGES820203


DISCARDED  
BY  
Reference Center

## NRE

Economic  
Research  
Service

United States  
Department  
of  
Agriculture





AD-33 Bookplate  
(1-68)

**NATIONAL**

**A  
G  
R  
I  
C  
U  
L  
T  
U  
R  
A  
L**



**LIBRARY**

1980 PESTICIDE USE ON SOYBEANS  
IN THE SOUTHEAST

by

Michael Hanthorn, Craig Osteen,  
Robert McDowell, and Larry Roberson

February 1982

ERS Staff Report No. AGES820203

U. S. DEPT. OF AGRICULTURE  
NATIONAL AGRICULTURAL LIBRARY

MAY 28 1986

CATALOGING = PREP.

Natural Resource Economics Division  
Economic Research Service  
U.S. Department of Agriculture  
Washington, D.C. 20250





1980 PESTICIDE USE ON SOYBEANS IN THE SOUTHEAST. By Michael Hanthorn, Craig Osteen, Robert McDowell, and Larry Roberson; Natural Resource Economics Division, Economic Research Service, U.S. Department of Agriculture, Washington, D.C. 20250; February 1982.

ERS Staff Report No. AGES820203

#### ABSTRACT

Farmers reported that 18.1 million pounds active ingredient (a.i.) of pesticides were applied to soybeans in the Southeast during 1980. This consisted of 11.3 million pounds (a.i.) of herbicides, 5.1 million pounds (a.i.) of insecticides, 1.4 million pounds (a.i.) of nematocides, and 234,000 pounds (a.i.) of fungicides. Pesticide acre-treatments totaled 16.4 million and consisted of 9.7 million with herbicides, 5.8 million with insecticides, 555,000 with nematocides, and 384,000 with fungicides. The primary herbicides were alachlor, bentazon, metribuzin, and trifluralin. The major insecticides were carbaryl and methomyl. Aldicarb and ethylene dibromide were the primary nematocides and benomyl was the major fungicide. Herbicides were applied primarily to control cocklebur and crabgrass infestations, while insecticides were mainly used for armyworm and corn earworm control. Anthracnose and pod and stem blight were the major diseases. Coefficients of variation were computed for acres of soybeans treated with specific pesticide materials.

Key words: Pesticides, herbicides, insecticides, nematocides, fungicides, active ingredient, acres treated, acre-treatments, application rates, primary target pests, soybeans, and Southeast.

\* \* \* \* \*

\* This paper was prepared for limited distribution to the research community outside the U.S. Department of Agriculture. Use of product names in this report is for identification only, and does not imply endorsement by the U.S. Department of Agriculture.

\* \* \* \* \*

#### ACKNOWLEDGMENTS

The 1980 Soybean Pesticide Use Survey was conducted by the Statistics Division of the Economics and Statistics Service. Herman W. Delvo provided guidance and made valuable suggestions during this effort. The report was reviewed by Walter Ferguson and Edward O. Fryar, Jr. A special thanks is extended to Andrea Lunsford for typing the preliminary and final manuscript drafts.

#### AUTHORS

Hanthorn, Osteen, and McDowell are with the Economic Research Service. Roberson is with the Statistical Reporting Service.

## PREFACE

This report presents data for pesticides applied to soybeans in the Southeast during 1980. Pesticide use data for the major producing States not included in the Southeast and for all major producing regions are available in the following ERS Staff Reports:

- "1980 Pesticide Use on Soybeans in the North Central States"
- "1980 Pesticide Use on Soybeans in the Mississippi Valley"
- "1980 Pesticide Use on Soybeans in the Major Producing States".

## CONTENTS

|                                   | <u>Page</u> |
|-----------------------------------|-------------|
| INTRODUCTION .....                | 1           |
| METHODOLOGY AND TERMINOLOGY ..... | 1           |
| RELIABILITY OF ESTIMATES .....    | 2           |
| SOUTHEAST .....                   | 3           |
| Description .....                 | 3           |
| Trends in Pesticide Use .....     | 3           |
| Pesticide Use .....               | 7           |
| ALABAMA .....                     | 14          |
| GEORGIA .....                     | 16          |
| NORTH CAROLINA .....              | 18          |
| SOUTH CAROLINA .....              | 21          |
| REFERENCES .....                  | 24          |
| APPENDIX TABLES .....             | 25          |



## INTRODUCTION

This report presents pesticide use data for soybeans grown in the Southeast during 1980. The data include usage patterns and quantities of specific herbicides, insecticides, nematicides, and fungicides applied to soybeans. This information should be useful to policymakers, academic institutions, government agencies, and private and commercial entities in evaluating the impacts of regulatory actions on specific pesticides, conducting economic analyses of pesticide use, developing more effective pest management programs, and conducting pesticide market analyses.

## METHODOLOGY AND TERMINOLOGY

The Economics and Statistics Service collected pesticide use data as part of the 1980 Soybean Objective Yield Survey. A total of 1,915 farmers were personally interviewed by enumerators in the 17 major soybean producing States. In the Southeast, 310 farmers were interviewed and the sample size by State was as follows: Alabama, 80; Georgia, 75; North Carolina, 80; and South Carolina, 75.

Sample fields for each State were randomly selected from farmers who reported through the June Enumerative Survey that they had planted or intended to plant soybeans in 1980. Each soybean acre in a State had an equal probability of being selected. Consequently, the probability of a field being chosen was directly correlated to its size.

Several terms pertinent to this report are defined as follows. An "active ingredient" (a.i.) is that portion of a pesticide material that provides the control activity. "Acres treated" are the number of acres receiving one or more applications of a specific pesticide during the growing season. Acres treated with different pesticide materials cannot be summed because more than

one material may have been applied on a given acre during the growing season. Therefore, the addition of these numbers would result in multiple counting.

"Acre-treatments" are the number of acres treated with a pesticide material multiplied by the number of applications made during the growing season. Acre-treatments are summed for each material at the State and regional level. "Pesticide mixes" are two or more pesticide materials that are premixed during formulation or tank-mixed at the time of application.

Pesticide application rates vary as a result of weather conditions, soil type, weed spectrum, insect species, and disease type. Also, the method of application influences the amount of a material used per acre. Herbicide and foliar insecticide application rates are generally expressed as broadcast rates. The amount of a material applied on an acre in either a band, in-furrow, or spot application is generally one-fourth to one-third the amount applied in a broadcast application. The application rate listed for each material in this report is an aggregation of band, broadcast, in-furrow, and spot applications.

#### RELIABILITY OF ESTIMATES

Estimates based upon sample surveys have varying degrees of statistical reliability. Confidence in data depends upon sample size, sampling methods, and the variability of the responses. To provide the user of the data with some indication of the reliability of the estimates, coefficients of variation (CV's) are presented in Appendix Table 1. The CV is a measure of relative variation (expressed in percentage terms) and can be used to indicate the degree of confidence a user can place in the estimate. The smaller the CV, the more reliable the estimate.

In simplest terms, it can be said there is 95 percent confidence that the sample represents the true population and that the true value for the population



lies within an interval defined as  $\pm 2$  CV's times the estimated value. For example, with a CV of 10 percent and an estimate of 40, the interval would be 32 to 48. However, there is also a 5 percent chance that the true value does not fall within the interval as defined above because the sample is not representative of the population.

CV's were calculated only for acres treated with specific pesticides. The estimates of acres treated are expected to have greater variation than other data reported. Consequently, for most other information included in this report, the level of reliability should be equal to or greater than that reported for acres treated.

## SOUTHEAST

### Description

The Southeast includes Alabama, Georgia, North Carolina, and South Carolina (Figure 1). About 12 percent of the U.S. soybean acreage (8.5 million acres) was planted in this region, from which 6 percent of the national crop (116 million bushels) was produced (Table 1). The farm value of soybeans grown in this region during 1980 was \$886 million.

### Trends in Pesticide Use

There was a substantial increase in the number of acres planted to soybeans and treated with pesticides in the Southeast between 1972 and 1980. Acres planted more than doubled from 3.9 to 8.5 million, while the percent of planted acres treated with herbicides and insecticides increased 65 and 370 percent, respectively (Table 2). Approximately three-fourths of the planted acres were treated with herbicides in 1980, while 47 percent were treated with insecticides. Planted acres treated with nematocides totaled 531,000 (6 percent) and fungicide treated acres totaled 208,000 (2 percent).

Figure 1. States included in the 1980 Soybean Pesticide Use Survey






Table 1. Soybean acreage planted and harvested, production, and value in the Southeast, 1980

| States                   | : Total acres a/ :       |             | : Total : Total                  |                                  |
|--------------------------|--------------------------|-------------|----------------------------------|----------------------------------|
|                          | : Planted                | : Harvested | : production a/                  | : value b/                       |
|                          | <u>-----Million-----</u> |             | <u>Million</u><br><u>bushels</u> | <u>Million</u><br><u>dollars</u> |
| Alabama                  | 2.3                      | 2.1         | 32                               | 239                              |
| Georgia                  | 2.5                      | 2.2         | 26                               | 194                              |
| North Carolina           | 2.0                      | 1.9         | 36                               | 280                              |
| South Carolina           | 1.7                      | 1.6         | 22                               | 173                              |
| Region                   | 8.5                      | 7.8         | 116                              | 886                              |
| U.S. Total               | 70.1                     | 67.9        | 1,817                            | 13,825                           |
| Percent of<br>U.S. Total | 12                       | 11          | 6                                | 6                                |

a/ "Crop Production-1980 Annual Summary", USDA, ESS, Crop Reporting Board, CrPr 2-1(81), January 14, 1981.

b/ "Field Crops-Production, Disposition, Value 1979-80", USDA, ESS, Crop Reporting Board, CrPr 1(81), April 1981.

Table 2. Soybean acreage planted and treated for weed, insect, nematode, and disease control in the Southeast, 1972 and 1980

| States         | Planted                          |         | Treated acres a/ |         |              |         | Percent of planted acres treated |       |              |      |
|----------------|----------------------------------|---------|------------------|---------|--------------|---------|----------------------------------|-------|--------------|------|
|                | acres                            |         | Herbicides       |         | Insecticides |         | Herbicides                       |       | Insecticides |      |
|                | 1972 b/                          | 1980 c/ | 1972 d/          | 1980 e/ | 1972 d/      | 1980 e/ | 1972:                            | 1980: | 1972:        | 1980 |
|                | -----Thousand----- Percent ----- |         |                  |         |              |         |                                  |       |              |      |
| Alabama        | 830                              | 2,250   | -                | 1,832   | -            | 547     | -                                | 81    | -            | 24   |
| Georgia        | 720                              | 2,450   | -                | 1,881   | -            | 1,517   | -                                | 77    | -            | 62   |
| North Carolina | 1,250                            | 2,030   | 535              | 1,421   | 126          | 821     | 43                               | 70    | 10           | 40   |
| South Carolina | 1,130                            | 1,700   | 567              | 1,298   | 108          | 1,086   | 50                               | 76    | 10           | 64   |
| Region         | 3,930                            | 8,430   | 1,102            | 6,432   | 234          | 3,971   | 46f/                             | 76    | 10f/         | 47   |

- Not surveyed in 1972.

a/ Nematicide and fungicide use was not reported in these States in 1972.

In 1980, the number of planted acres treated with nematicides in these States was as follows: Alabama, 161,000 (7 percent); Georgia, none reported; North Carolina, 87,000 (4 percent); and South Carolina, 283,000 (17 percent). Fungicide treated acres were as follows: Alabama, 32,000 (1 percent); Georgia, 71,000 (3 percent); North Carolina, 58,000 (3 percent); and South Carolina, 47,000 (3 percent).

b/ "Agricultural Statistics, 1974", U.S. Department of Agriculture.

c/ "Crop Production-1980 Annual Summary", USDA, ESS, Crop Reporting Board, CrPr 2-1(81), January 14, 1981.

d/ Herman W. Delvo, "1972 Soybean Objective Yield Survey", USDA, ERS, Farm Production Economics Division, 1972, (unpublished).

e/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

f/ Computed excluding planted acres for Alabama and Georgia.

## Pesticide Use

The major soybean weed, insect, and disease pests, as reported by farmers in the Southeast, are listed in Tables 3, 4, and 5, respectively. Although several pests may have been present at any given time and caused varying degrees of damage, farmers were asked to report what they perceived to be the primary target pest for each material applied to soybeans. In 1980, farmers reported that cocklebur was the primary target pest for 34 percent of the herbicide acre-treatments, crabgrass for 25 percent, and Johnsongrass for 8 percent (Table 3). Insecticide acre-treatments for corn earworm and armyworm control totaled 39 and 31 percent, respectively (Table 4). Two-thirds of the fungicide acre-treatments were made to control pod and stem blight and 23 percent were made for anthracnose control (Table 5).

About 18.1 million pounds (a.i.) of pesticides were applied to soybeans in 1980 (Table 6). Of these, 6.5 million were single material herbicides, 4.8 million were herbicide mixes, 3.9 million were single material insecticides, 1.2 million were insecticide mixes, 1.4 million were nematicides, and 234,000 were fungicides. Application rates for herbicides, applied alone and in mixes, were 0.9 and 2.2 pounds (a.i.) per acre-treatment, respectively. Insecticide application rates averaged 0.8 pound (a.i.) per acre-treatment for single materials and 1.6 pounds (a.i.) per acre-treatment for mixes. Nematicide and fungicide rates averaged 2.6 and 0.6 pounds (a.i.) per acre-treatment, respectively.

Farmers made 16.4 million pesticide acre-treatments, comprised of 7.5 million with single material herbicides, 2.2 million with herbicide mixes, 5 million with single material insecticides, 764,000 with insecticide mixes, 555,000 with nematicides, and 384,000 with fungicides.

About 2.8 million (37 percent) of the single material herbicide acre-treatments were trifluralin, while alachlor, bentazon, and metribuzin acre-treatments

Table 3. Percentage of soybean herbicide acre-treatments by primary weeds controlled as reported by farmers in the Southeast, 1980 a/

|                       | :  | :       | :  | : North        | : South | :                            |
|-----------------------|----|---------|----|----------------|---------|------------------------------|
| Weeds                 | :  | Alabama | :  | Georgia        | :       | Carolina : Carolina : Region |
|                       |    |         |    | <u>Percent</u> |         |                              |
| <u>Grasses</u>        |    |         |    |                |         |                              |
| Barnyardgrass         | 3  | -       | -  | -              | 1       |                              |
| Broadleaf signalgrass | -  | 4       | 5  | 4              | 3       |                              |
| Crabgrass             | 13 | 36      | 19 | 34             | 25      |                              |
| Johnsongrass          | 15 | 2       | 11 | 3              | 8       |                              |
| Panicum               | 1  | -       | 5  | 4              | 2       |                              |
| Other                 | 4  | 8       | 5  | 3              | 5       |                              |
| <u>Broadleaves</u>    |    |         |    |                |         |                              |
| Cocklebur             | 40 | 29      | 26 | 39             | 34      |                              |
| Morningglory          | 5  | -       | 10 | 2              | 4       |                              |
| Pigweed               | 3  | 6       | 5  | 6              | 5       |                              |
| Ragweed               | 3  | -       | 5  | -              | 2       |                              |
| Sicklepod             | 8  | 8       | 2  | 2              | 5       |                              |
| Smartweed             | 1  | -       | -  | -              | 1       |                              |
| Other                 | 3  | 6       | 5  | 2              | 4       |                              |
| Nutsedge              | 1  | 1       | 2  | 1              | 1       |                              |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.



Table 4. Percentage of soybean insecticide acre-treatments by primary insects controlled as reported by farmers in the Southeast, 1980 a/

| Insects                | : Alabama      | : Georgia | : North Carolina | : South Carolina | : Region |
|------------------------|----------------|-----------|------------------|------------------|----------|
|                        | <u>Percent</u> |           |                  |                  |          |
| Armyworm               | 33             | 42        | 15               | 23               | 31       |
| Bean leaf beetle       | -              | -         | -                | 8                | 2        |
| Cabbage looper         | -              | 10        | 19               | 3                | 7        |
| Corn earworm           | 48             | 34        | 35               | 42               | 39       |
| Cutworm                | -              | 6         | 6                | 1                | 4        |
| Grasshopper            | -              | 4         | -                | -                | 2        |
| Green cloverworm       | -              | -         | 9                | -                | 2        |
| Other beetles          | -              | -         | -                | 11               | 3        |
| Spider mite            | -              | -         | 9                | -                | 2        |
| Velvetbean caterpillar | -              | 2         | 3                | 8                | 4        |
| Other                  | 19             | 2         | 4                | 4                | 4        |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

Table 5. Percentage of soybean fungicide acre-treatments by primary diseases controlled as reported by farmers in the Southeast, 1980 a/

| Diseases            | : Alabama      | : Georgia | : North Carolina | : South Carolina | : Region |
|---------------------|----------------|-----------|------------------|------------------|----------|
|                     | <u>Percent</u> |           |                  |                  |          |
| Anthracnose         | -              | -         | 100              | -                | 23       |
| Leaf blight         | -              | 33        | -                | -                | 9        |
| Pod and stem blight | 100            | 67        | -                | 100              | 68       |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

Table 6. Usage patterns and quantities of specific pesticides applied to soybeans in the Southeast, 1980 a/

| Pesticides                   | : Acres <u>b/</u> | : Acre- <u>c/</u> | : Pounds of active ingredient |                 |
|------------------------------|-------------------|-------------------|-------------------------------|-----------------|
|                              | : treated         | : treatments:     | Total                         | : Per treatment |
| ----- <u>Thousand</u> -----  |                   |                   |                               |                 |
| HERBICIDES                   |                   |                   |                               |                 |
| <u>Single materials</u>      |                   |                   |                               |                 |
| Alachlor                     | 1,082             | 1,082             | 1,719                         | 1.6             |
| Bentazon                     | 1,085             | 1,176             | 765                           | .7              |
| Metribuzin                   | 896               | 896               | 481                           | .5              |
| Pendimethalin                | 262               | 262               | 252                           | 1.0             |
| Trifluralin                  | 2,713             | 2,769             | 2,000                         | .7              |
| Other                        | -                 | 1,297             | 1,243                         | 1.0             |
| Total                        | -                 | 7,482             | 6,460                         | .9              |
| <u>Tank-mix materials</u>    |                   |                   |                               |                 |
| Alachlor + linuron           | 263               | 263               | 593+221                       | 2.3+ .8         |
| Dinoseb + naptalam           | 587               | 587               | 428+839                       | .7+1.4          |
| Metribuzin + trifluralin     | 290               | 290               | 107+218                       | .4+1.8          |
| Other                        | -                 | 1,068             | 2,433                         | 2.3             |
| Total                        | -                 | 2,208             | 4,839                         | 2.2             |
| Total herbicides             | -                 | 9,690             | 11,299                        | 1.2             |
| INSECTICIDES                 |                   |                   |                               |                 |
| <u>Single materials</u>      |                   |                   |                               |                 |
| Carbaryl                     | 877               | 1,282             | 1,358                         | 1.1             |
| Methomyl                     | 1,716             | 2,542             | 1,198                         | .5              |
| Methyl parathion             | 248               | 248               | 190                           | .8              |
| Parathion                    | 144               | 208               | 187                           | .9              |
| Toxaphene                    | 342               | 484               | 889                           | 1.8             |
| Other                        | -                 | 273               | 91                            | .3              |
| Total                        | -                 | 5,037             | 3,913                         | .8              |
| <u>Tank-mix materials</u>    |                   |                   |                               |                 |
| Methyl parathion + toxaphene | 150               | 324               | 372+425                       | 1.1+1.3         |
| Other                        | -                 | 440               | 400                           | .9              |
| Total                        | -                 | 764               | 1,197                         | 1.6             |
| Total insecticides           | -                 | 5,801             | 5,110                         | .9              |
| NEMATICIDES                  |                   |                   |                               |                 |
| Aldicarb                     | 214               | 214               | 279                           | 1.3             |
| Carbofuran                   | 114               | 114               | 91                            | .8              |
| Ethoprop                     | 53                | 53                | 117                           | 2.2             |
| Ethylene dibromide           | 142               | 142               | 783                           | 5.5             |
| Other                        | -                 | 32                | 154                           | 4.8             |
| Total                        | -                 | 555               | 1,424                         | 2.6             |

-- continued

Table 6. Usage patterns and quantities of specific pesticides applied to soybeans in the Southeast, 1980 a/ -- continued

| Pesticides                  | : Acres <u>b/</u> | : Acre- <u>c/</u> | :Pounds of active ingredient |                 |
|-----------------------------|-------------------|-------------------|------------------------------|-----------------|
|                             | : treated         | :treatments:      | Total                        | : Per treatment |
| ----- <u>Thousand</u> ----- |                   |                   |                              |                 |
| FUNGICIDES                  |                   |                   |                              |                 |
| Benomyl                     | 150               | 297               | 83                           | .3              |
| Benomyl + other             | -                 | 87                | 90+61                        | 1.0+ .7         |
| Total                       | -                 | 384               | 234                          | .6              |
| TOTAL PESTICIDES            | -                 | 16,430            | 18,067                       | 1.1             |

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ Data in this column for "other" and "total" were not reported because two or more materials may have been used on the same acre resulting in multiple counting.

c/ Most farmers applied herbicides, insecticides, nematocides, and fungicides one time during the growing season. The average number of applications per season for each material can be determined by dividing acre-treatments (column 2) by acres treated (column 1).



totalled 3.2 million (42 percent). Dinoseb plus naptalam acre-treatments accounted for 587,000 (27 percent) of the herbicide mix acre-treatments. Also, one-fourth (553,000) of these acre-treatments were made with alachlor plus linuron and metribuzin plus trifluralin.

Crabgrass control accounted for 35 percent of the alachlor acre-treatments and 50 percent of the trifluralin acre-treatments (Appendix Table 2). About 12 percent of the alachlor acre-treatments, 91 percent of the bentazon acre-treatments, 41 percent of the metribuzin acre-treatments, and 15 percent of the trifluralin acre-treatments were made to control cocklebur infestations. Approximately 18 percent of the metribuzin acre-treatments were made to control morningglory infestations and 15 percent of the trifluralin acre-treatments were made for Johnsongrass control.

Methomyl accounted for 2.5 million (45 percent) of the single material insecticide acre-treatments (Table 6). Also, 1.3 million (23 percent) were carbaryl. Methyl parathion plus toxaphene accounted for 324,000 (42 percent) of the insecticide mix acre-treatments. Carbaryl acre-treatments totaled 37 and 30 percent, respectively, for suppression of armyworm and corn earworm infestations (Appendix Table 3). Nearly one-half of the methomyl acre-treatments were made for corn earworm control and one-fifth were made to suppress armyworm infestations. Armyworm control accounted for 43 percent of the methyl parathion acre-treatments and velvetbean caterpillar control accounted for 10 percent.

Aldicarb, carbofuran, and ethylene dibromide accounted for 470,000 (85 percent) of the nematicide acre-treatments (Table 6). Three-fourths (297,000) of the fungicide acre-treatments were made with benomyl and the remainder (87,000) were benomyl plus other pesticides. About 88 percent of the benomyl acre-treatments were made to control pod and stem blight and all of the benomyl mixed with other pesticides was used for anthracnose control (Appendix Table 4).

## ALABAMA

In 1980, Alabama farmers planted 2.3 million acres of soybeans, of which 1.8 million were treated with herbicides, 547,000 were treated with insecticides, 161,000 were treated with nematicides, and 32,000 were treated with fungicides (Table 2). Approximately 3.8 million pounds (a.i.) of pesticides were applied to soybeans, which included 1.5 million pounds of single material herbicides, 1 million pounds of herbicide mixes, 425,000 pounds of single material insecticides, 450,000 pounds of insecticide mixes, 305,000 pounds of nematicides, and 24,000 pounds of fungicides (Table 7). Application rates for herbicides averaged 0.7 pound (a.i.) per acre-treatment for single materials and 1.6 pounds (a.i.) per acre-treatment for mixes. Insecticide rates averaged 0.7 pound (a.i.) per acre-treatment for single materials and 2 pounds (a.i.) per acre-treatment for mixes. Nematicide and fungicide rates averaged 1.9 and 0.3 pounds (a.i.) per acre-treatment, respectively.

Pesticide acre-treatments totaled 4.1 million, comprised of 2.4 million with single material herbicides, 644,000 with herbicide mixes, 580,000 with single material insecticides, 225,000 with insecticide mixes, 160,000 with nematicides, and 96,000 with fungicides.

One-third (788,000) of the single material herbicide acre-treatments were trifluralin, one-fourth (583,000) were bentazon, and 15 percent (354,000) were metribuzin. One-fourth (161,000) of the herbicide mix acre-treatments were metribuzin plus trifluralin and 10 percent (64,000) were acifluorfen plus bentazon. Cocklebur, crabgrass, and sicklepod control each accounted for one-sixth of the alachlor acre-treatments, while one-half were made for barnyardgrass control (Appendix Table 2). Cocklebur control accounted for 94 percent of the bentazon acre-treatments. Metribuzin acre-treatments totaled 37 percent for cocklebur control and 36 percent for morningglory control. Nearly one-third

Table 7. Usage patterns and quantities of specific pesticides applied to soybeans in Alabama, 1980 a/

| Pesticides                   | : Acres <u>b/</u><br>: treated | : Acre- <u>c/</u><br>: treatments: | : Pounds of active ingredient<br>Total : Per treatment |         |
|------------------------------|--------------------------------|------------------------------------|--|---------|
|                              |                                |                                    |  |         |
| ----- <u>Thousand</u> -----  |                                |                                    |  |         |
| HERBICIDES                   |                                |                                    |  |         |
| <u>Single materials</u>      |                                |                                    |  |         |
| Alachlor                     | 192                            | 192                                | 206  | 1.1     |
| Bentazon                     | 551                            | 583                                | 293  | .5      |
| Metribuzin                   | 354                            | 354                                | 216  | .6      |
| Trifluralin                  | 755                            | 788                                | 522  | .7      |
| Other                        | -                              | 448                                | 301  | .7      |
| Total                        | -                              | 2,365                              | 1,538  | .7      |
| <u>Tank-mix materials</u>    |                                |                                    |  |         |
| Acifluorfen + bentazon       | 64                             | 64                                 | 23+48  | .4+ .8  |
| Metribuzin + trifluralin     | 161                            | 161                                | 45+109   | .3+ .7  |
| Other                        | -                              | 419                                | 808  | 1.9     |
| Total                        | -                              | 644                                | 1,033  | 1.6     |
| Total herbicides             | -                              | 3,009                              | 2,571  | .9      |
| INSECTICIDES                 |                                |                                    |  |         |
| <u>Single materials</u>      |                                |                                    |  |         |
| Methomyl                     | 232                            | 264                                | 138  | .5      |
| Parathion                    | 96                             | 161                                | 161  | 1.0     |
| Toxaphene                    | 64                             | 64                                 | 64   | 1.0     |
| Other                        | -                              | 91                                 | 62   | .7      |
| Total                        | -                              | 580                                | 425  | .7      |
| <u>Tank-mix materials</u>    |                                |                                    |  |         |
| Methomyl + toxaphene         | 32                             | 96                                 | 48+48  | .5+ .5  |
| Methyl parathion + toxaphene | 32                             | 129                                | 177+177  | 1.4+1.4 |
| Total                        | -                              | 225                                | 450  | 2.0     |
| Total insecticides           | -                              | 805                                | 875  | 1.1     |
| NEMATICIDES                  |                                |                                    |  |         |
| Aldicarb                     | 96                             | 96                                 | 145  | 1.5     |
| Other                        | -                              | 64                                 | 160  | 2.5     |
| Total                        | -                              | 160                                | 305  | 1.9     |
| FUNGICIDES                   |                                |                                    |  |         |
| Total <u>d/</u>              | -                              | 96                                 | 24   | .3      |
| TOTAL PESTICIDES             | -                              | 4,070                              | 3,775  | .9      |

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ Data in this column for "other" and "total" were not reported because two or more materials may have been used on the same acre resulting in multiple counting.

c/ Most farmers applied herbicides, insecticides, nematicides, and fungicides one time during the growing season. The average number of applications per season for each material can be determined by dividing acre-treatments (column 2) by acres treated (column 1).

d/ Specific materials not listed because of a limited number of observations.

of the trifluralin acre-treatments were made to control crabgrass infestations, while 37 percent were made for Johnsongrass control.

Methomyl accounted for 264,000 (46 percent) of the single material insecticide acre-treatments, while parathion and toxaphene acre-treatments totaled 161,000 (28 percent) and 64,000 (11 percent), respectively (Table 7). About 129,000 (57 percent) of the insecticide mix acre-treatments were made with methyl parathion plus toxaphene and 96,000 (43 percent) were methomyl plus toxaphene. Two-thirds of the methomyl acre-treatments were made for armyworm control and one-fourth were made to suppress corn earworm infestations (Appendix Table 3).

Aldicarb accounted for 96,000 (60 percent) of the nematicide acre-treatments (Table 7). Fungicide acre-treatments totaled 96,000 and were made to control pod and stem blight (Table 5).

#### GEORGIA

Acres planted to soybeans totaled 2.5 million in Georgia during 1980, of which 1.9 million were treated with herbicides, 1.5 million were treated with insecticides, and 71,000 were treated with fungicides (Table 2). A total of 6.2 million pounds (a.i.) of pesticides were applied to soybeans, which consisted of 1.9 million pounds of single material herbicides, 2 million pounds of herbicide mixes, 1.7 million pounds of single material insecticides, 558,000 pounds of insecticide mixes, and 36,000 pounds of fungicides (Table 8). Application rates for herbicides averaged 0.9 pound (a.i.) per acre-treatment for single materials and 3 pounds (a.i.) per acre-treatment for mixes. Rates for insecticides, applied alone and in mixes, averaged 0.9 and 1.3 pounds (a.i.) per acre-treatment, respectively. Fungicide rates averaged 0.3 pound (a.i.) per acre-treatment.

Pesticide acre-treatments totaled 5.4 million and were comprised of 2.2



Table 8. Usage patterns and quantities of specific pesticides applied to soybeans in Georgia, 1980 a/

| Pesticides                   | : Acres <u>b/</u> | : Acre- <u>c/</u> | :Pounds of active ingredient |               |
|------------------------------|-------------------|-------------------|------------------------------|---------------|
|                              | : treated         | :treatments:      | Total                        | Per treatment |
| ----- <u>Thousand</u> -----  |                   |                   |                              |               |
| HERBICIDES                   |                   |                   |                              |               |
| <u>Single materials</u>      |                   |                   |                              |               |
| Alachlor                     | 284               | 284               | 333                          | 1.2           |
| Bentazon                     | 192               | 192               | 140                          | .7            |
| Metribuzin                   | 379               | 379               | 189                          | .5            |
| Oryzalin                     | 134               | 134               | 117                          | .9            |
| Pendimethalin                | 142               | 142               | 160                          | 1.1           |
| Trifluralin                  | 849               | 849               | 715                          | .8            |
| Other                        | -                 | 214               | 246                          | 1.1           |
| Total                        | -                 | 2,194             | 1,900                        | .9            |
| <u>Tank-mix materials</u>    |                   |                   |                              |               |
| Dinoseb + naptalam           | 426               | 426               | 362+713                      | .9+1.7        |
| Other                        | -                 | 249               | 959                          | 3.9           |
| Total                        | -                 | 675               | 2,034                        | 3.0           |
| Total herbicides             | -                 | 2,869             | 3,934                        | 1.4           |
| INSECTICIDES                 |                   |                   |                              |               |
| <u>Single materials</u>      |                   |                   |                              |               |
| Carbaryl                     | 71                | 107               | 96                           | .9            |
| Methomyl                     | 600               | 1,059             | 546                          | .5            |
| Methyl parathion             | 142               | 142               | 142                          | 1.0           |
| Permethrin                   | 178               | 178               | 18                           | .1            |
| Toxaphene                    | 278               | 420               | 824                          | 2.0           |
| Other                        | -                 | 36                | 29                           | .8            |
| Total                        | -                 | 1,942             | 1,655                        | .9            |
| <u>Tank-mix materials</u>    |                   |                   |                              |               |
| Methyl parathion + toxaphene | 71                | 124               | 142+142                      | 1.1+1.1       |
| Other                        | -                 | 320               | 274                          | .9            |
| Total                        | -                 | 444               | 558                          | 1.3           |
| Total insecticides           | -                 | 2,386             | 2,213                        | .9            |
| FUNGICIDES                   |                   |                   |                              |               |
| Benomyl                      | 71                | 107               | 36                           | .3            |
| TOTAL PESTICIDES             | -                 | 5,362             | 6,183                        | 1.2           |

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ Data in this column for "other" and "total" were not reported because two or more materials may have been used on the same acre resulting in multiple counting.

c/ Most farmers applied herbicides, insecticides, and fungicides one time during the growing season. The average number of applications per season for each material can be determined by dividing acre-treatments (column 2) by acres treated (column 1).

million with single material herbicides, 675,000 with herbicide mixes, 1.9 million with single material insecticides, 444,000 with insecticide mixes, and 107,000 with fungicides.

Of the 2.2 million single material herbicide acre-treatments, 849,000 (39 percent) were trifluralin, 379,000 (17 percent) were metribuzin, and 284,000 (13 percent) were alachlor. Nearly two-thirds (426,000) of the herbicide mix acre-treatments were dinoseb plus naptalam. Three-fourths of the alachlor and 62 percent of the trifluralin acre-treatments were made to control crabgrass infestations (Appendix Table 2). All of the bentazon acre-treatments and 56 percent of the metribuzin acre-treatments were made for cocklebur control.

Methomyl accounted for 1.1 million (55 percent) of the single material insecticide acre-treatments and toxaphene acre-treatments totaled 420,000 (22 percent) (Table 8). Methyl parathion plus toxaphene accounted for 124,000 (28 percent) of the insecticide mix acre-treatments. All of the carbaryl, 75 percent of the methyl parathion, and 20 percent of the methomyl acre-treatments were made for armyworm control (Appendix Table 3). Corn earworm control accounted for 57 percent of the methomyl acre-treatments.

Benomyl acre-treatments totaled 107,000 (Table 8). Two-thirds were made for pod and stem blight control and the remainder were made for leaf blight control (Appendix Table 4).

#### NORTH CAROLINA

About 2 million acres of soybeans were planted in North Carolina during the 1980 growing season (Table 2). Of these, 1.4 million were treated with herbicides, 821,000 were treated with insecticides, 87,000 were treated with nematicides, and 58,000 were treated with fungicides. A total of 4 million pounds (a.i.) of pesticides were applied to soybeans, consisting of 2 million

pounds of single material herbicides, 853,000 pounds of herbicide mixes, 893,000 pounds of insecticides, 107,000 pounds of nematocides, and 151,000 pounds of fungicides (Table 9). An estimated 1.5 pounds (a.i.) of single material herbicides and 2 pounds (a.i.) of herbicide mixes were applied per acre-treatment. Insecticides were applied at an average rate of 1 pound (a.i.) per acre-treatment. Nematicide and fungicide rates averaged 1.2 and 1.7 pounds (a.i.) per acre-treatment, respectively.

Pesticide acre-treatments totaled 2.9 million and consisted of 1.4 million with single material herbicides, 435,000 with herbicide mixes, 937,000 with insecticides, 87,000 with nematocides, and 87,000 with fungicides.

One-third (464,000) of the single material herbicide acre-treatments were alachlor, 174,000 (13 percent) were bentazon, and 232,000 (17 percent) were trifluralin. One-third (145,000) of the herbicide mix acre-treatments were made with alachlor plus linuron, while dinoseb plus naptalam and metribuzin plus trifluralin acre-treatments each totaled 58,000 (13 percent). Crabgrass and morningglory control each accounted for 19 percent of the alachlor acre-treatments, while 38 percent were made to reduce broadleaf signalgrass, cocklebur, and Johnsongrass infestations (Appendix Table 2). Cocklebur control accounted for two-thirds of the bentazon acre-treatments. Crabgrass, morningglory, pigweed, and ragweed control each accounted for 25 percent of the metribuzin acre-treatments. About 63 percent of the trifluralin acre-treatments were made to suppress crabgrass infestations.

One-half (464,000) of the insecticide acre-treatments were methomyl and 47 percent (444,000) were carbaryl (Table 9). Carbaryl acre-treatments totaled 35 percent for corn earworm control, 20 percent for armyworm control, 19 percent for green cloverworm control, and 26 percent for cabbage looper and cutworm control (Appendix Table 3). Corn earworm was the primary target pest for 38

Table 9. Usage patterns and quantities of specific pesticides applied to soybeans in North Carolina, 1980 a/

| Pesticides                  | : Acres <u>b/</u><br>: treated | : Acre- <u>c/</u><br>: treatments: | : Pounds of active ingredient |               |
|-----------------------------|--------------------------------|------------------------------------|-------------------------------|---------------|
|                             |                                |                                    | Total                         | Per treatment |
| ----- <u>Thousand</u> ----- |                                |                                    |                               |               |
| HERBICIDES                  |                                |                                    |                               |               |
| <u>Single materials</u>     |                                |                                    |                               |               |
| Alachlor                    | 464                            | 464                                | 1,087                         | 2.3           |
| Bentazon                    | 174                            | 174                                | 145                           | .8            |
| Metribuzin                  | 116                            | 116                                | 58                            | .5            |
| Trifluralin                 | 232                            | 232                                | 167                           | .7            |
| Other                       | -                              | 377                                | 537                           | 1.4           |
| Total                       | -                              | 1,363                              | 1,994                         | 1.5           |
| <u>Tank-mix materials</u>   |                                |                                    |                               |               |
| Alachlor + linuron          | 145                            | 145                                | 262+80                        | 1.8+ .6       |
| Dinoseb + naptalam          | 58                             | 58                                 | 23+43                         | .4+ .7        |
| Metribuzin + trifluralin    | 58                             | 58                                 | 36+44                         | .6+ .8        |
| Other                       | -                              | 174                                | 365                           | 2.1           |
| Total                       | -                              | 435                                | 853                           | 2.0           |
| Total herbicides            | -                              | 1,798                              | 2,847                         | 1.6           |
| INSECTICIDES                |                                |                                    |                               |               |
| Carbaryl                    | 357                            | 444                                | 670                           | 1.5           |
| Methomyl                    | 435                            | 464                                | 211                           | .5            |
| Other                       | -                              | 29                                 | 12                            | .4            |
| Total                       | -                              | 937                                | 893                           | 1.0           |
| NEMATICIDES                 |                                |                                    |                               |               |
| Carbofuran                  | 58                             | 58                                 | 61                            | 1.1           |
| Other                       | -                              | 29                                 | 46                            | 1.6           |
| Total                       | -                              | 87                                 | 107                           | 1.2           |
| FUNGICIDES                  |                                |                                    |                               |               |
| Benomyl + other             | -                              | 87                                 | 151                           | 1.7           |
| TOTAL PESTICIDES            | -                              | 2,909                              | 3,998                         | 1.4           |

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ Data in this column for "other" and "total" were not reported because two or more materials may have been used on the same acre resulting in multiple counting.

c/ Most farmers applied herbicides, insecticides, nematocides, and fungicides one time during the growing season. The average number of applications per season for each material can be determined by dividing acre-treatments (column 2) by acres treated (column 1).



percent of the methomyl acre-treatments, while 25 percent were made for cabbage looper control and 19 percent were made to suppress spider mite infestations.

Two-thirds (58,000) of the nematicide acre-treatments were made with carbofuran (Table 9). Benomyl was the primary fungicide that farmers reported using in 1980. It was applied in combination with other fungicides in 87,000 acre-treatments and was used for anthracnose control (Appendix Table 4).

#### SOUTH CAROLINA

Farmers planted 1.7 million acres to soybeans in South Carolina during 1980 and treated 1.3 million with herbicides, 1.1 million with insecticides, 283,000 with nematicides, and 47,000 with fungicides (Table 2). Approximately 4.1 million pounds (a.i.) of pesticides were applied to soybeans, of which 1 million were single material herbicides, 920,000 were herbicide mixes, 1.1 million were insecticides, 1 million were nematicides, and 24,000 were fungicides (Table 10). Herbicide application rates averaged 0.7 pound (a.i.) per acre-treatment for single materials and 2 pounds (a.i.) per acre-treatment for mixes. Insecticide application rates averaged 0.7 pound (a.i.) per acre-treatment, nematicide rates averaged 3.3 pounds (a.i.) per acre-treatment, and fungicide rates averaged 0.3 pound (a.i.) per acre-treatment.

Of the 4.1 million pesticide acre-treatments, 1.6 million were single material herbicides, 455,000 were herbicide mixes, 1.7 million were insecticides, 308,000 were nematicides, and 94,000 were fungicides.

About 900,000 (58 percent) of the single material herbicide acre-treatments were trifluralin, 227,000 (15 percent) were bentazon, and 142,000 (9 percent) were alachlor. One-fourth (118,000) of the herbicide mix acre-treatments were made with alachlor plus linuron, 17 percent (77,000) were made with dinoseb plus naptalam, and 16 percent (71,000) were made with metribuzin plus trifluralin.



Table 10. Usage patterns and quantities of specific pesticides applied to soybeans in South Carolina, 1980 a/

| Pesticides                  | : Acres <u>b/</u> | : Acre- <u>c/</u> | :Pounds of active ingredient |                 |
|-----------------------------|-------------------|-------------------|------------------------------|-----------------|
|                             | : treated         | :treatments:      | Total                        | : Per treatment |
| ----- <u>Thousand</u> ----- |                   |                   |                              |                 |
| HERBICIDES                  |                   |                   |                              |                 |
| <u>Single materials</u>     |                   |                   |                              |                 |
| Alachlor                    | 142               | 142               | 93                           | 0.7             |
| Bentazon                    | 168               | 227               | 187                          | .8              |
| Linuron                     | 71                | 71                | 24                           | .3              |
| Trifluralin                 | 877               | 900               | 596                          | .7              |
| Other                       | -                 | 221               | 128                          | .6              |
| Total                       | -                 | 1,561             | 1,028                        | .7              |
| <u>Tank-mix materials</u>   |                   |                   |                              |                 |
| Alachlor + linuron          | 118               | 118               | 331+142                      | 2.8+1.2         |
| Dinoseb + naptalam          | 77                | 77                | 35+67                        | .5+ .9          |
| Metribuzin + trifluralin    | 71                | 71                | 27+65                        | .4+ .9          |
| Other                       | -                 | 189               | 253                          | 1.3             |
| Total                       | -                 | 455               | 920                          | 2.0             |
| Total herbicides            | -                 | 2,016             | 1,948                        | 1.0             |
| INSECTICIDES                |                   |                   |                              |                 |
| Carbaryl                    | 449               | 732               | 592                          | .8              |
| Methomyl                    | 449               | 756               | 303                          | .4              |
| Other                       | -                 | 188               | 233                          | 1.2             |
| Total                       | -                 | 1,676             | 1,128                        | .7              |
| NEMATICIDES                 |                   |                   |                              |                 |
| Aldicarb                    | 118               | 118               | 135                          | 1.1             |
| Ethylene dibromide          | 142               | 142               | 783                          | 5.5             |
| Other                       | -                 | 48                | 95                           | 2.0             |
| Total                       | -                 | 308               | 1,013                        | 3.3             |
| FUNGICIDES                  |                   |                   |                              |                 |
| Benomyl                     | 47                | 94                | 24                           | .3              |
| TOTAL PESTICIDES            | -                 | 4,094             | 4,113                        | 1.0             |

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ Data in this column for "other" and "total" were not reported because two or more materials may have been used on the same acre resulting in multiple counting.

c/ Most farmers applied herbicides, insecticides, nematocides, and fungicides one time during the growing season. The average number of applications per season for each material can be determined by dividing acre-treatments (column 2) by acres treated (column 1).

Crabgrass and panicum control each accounted for one-third of the alachlor acre-treatments (Appendix Table 2). The remaining third was made for Johnson-grass and nutsedge control. Cocklebur was the primary target pest for 90 percent of the bentazon, 50 percent of the metribuzin, and 29 percent of the trifluralin acre-treatments. One-half of the metribuzin acre-treatments were made to control sicklepod infestations and 52 percent of the trifluralin acre-treatments were made for crabgrass control.

Carbaryl accounted for 732,000 (44 percent) of the insecticide acre-treatments and methomyl acre-treatments totaled 756,000 (45 percent) (Table 10). One-third of the carbaryl acre-treatments and one-half of the methomyl acre-treatments were made to suppress corn earworm infestations (Appendix Table 3). Armyworm control accounted for 39 percent of the carbaryl and 13 percent of the methomyl acre-treatments. Velvetbean caterpillar was the primary target pest for one-half of the methyl parathion acre-treatments and one-sixth of the carbaryl acre-treatments.

Aldicarb and ethylene dibromide accounted for 118,000 (38 percent) and 142,000 (46 percent) of the nematocide acre-treatments, respectively (Table 10). Benomyl acre-treatments totaled 94,000 and were made for pod and stem blight control (Appendix Table 4).

REFERENCES

1. Delvo, Herman W., "1972 Soybean Objective Yield Survey", USDA, ERS, Farm Production Economics Division, 1972, (unpublished).
2. U.S. Department of Agriculture, "Agricultural Statistics, 1974".
3. USDA, ESS, Crop Reporting Board, "Crop Production-1980 Annual Summary", CrPr 2-1(81), January 14, 1981.
4. USDA, ESS, Crop Reporting Board, "Field Crops-Production, Disposition, Value 1979-80", CrPr 1(81), April 1981.

Appendix Table 1. Coefficients of variation for acres of soybeans treated with specific pesticides in the Southeast, 1980 a/ b/

|                                 |                            |         |           |           |    |          |   |          |   |        |
|---------------------------------|----------------------------|---------|-----------|-----------|----|----------|---|----------|---|--------|
|                                 | :                          | :       | :         | North     | :  | South    | : |          |   |        |
| Pesticides                      | :                          | Alabama | :         | Georgia   | :  | Carolina | : | Carolina | : | Region |
|                                 | ----- <u>Percent</u> ----- |         |           |           |    |          |   |          |   |        |
| HERBICIDES                      |                            |         |           |           |    |          |   |          |   |        |
| <u>Single materials</u>         |                            |         |           |           |    |          |   |          |   |        |
| Alachlor                        | 39                         | 33      | 22        | 39        | 16 |          |   |          |   |        |
| Bentazon                        | 21                         | 41      | 39        | 36        | 15 |          |   |          |   |        |
| Metribuzin                      | 28                         | 28      | 49        | 70        | 18 |          |   |          |   |        |
| Pendimethalin                   | 57                         | 49      | -         | <u>c/</u> | 35 |          |   |          |   |        |
| Trifluralin                     | 17                         | 17      | 34        | 11        | 8  |          |   |          |   |        |
| <u>Tank-mix materials</u>       |                            |         |           |           |    |          |   |          |   |        |
| Alachlor + linuron              | -                          | -       | 43        | 43        | 31 |          |   |          |   |        |
| Dinoseb + naptalam              | <u>c/</u>                  | 26      | 70        | 53        | 22 |          |   |          |   |        |
| Metribuzin<br>+ trifluralin     | 43                         | -       | 70        | 57        | 31 |          |   |          |   |        |
| INSECTICIDES                    |                            |         |           |           |    |          |   |          |   |        |
| <u>Single materials</u>         |                            |         |           |           |    |          |   |          |   |        |
| Carbaryl                        | -                          | 70      | 25        | 20        | 15 |          |   |          |   |        |
| Methomyl                        | 29                         | 17      | 19        | 19        | 10 |          |   |          |   |        |
| Methyl parathion                | 71                         | 49      | -         | 70        | 35 |          |   |          |   |        |
| Parathion                       | 57                         | -       | -         | 70        | 45 |          |   |          |   |        |
| Toxaphene                       | 53                         | 32      | -         | -         | 28 |          |   |          |   |        |
| <u>Tank-mix materials</u>       |                            |         |           |           |    |          |   |          |   |        |
| Methyl parathion<br>+ toxaphene | 70                         | 70      | -         | 70        | 41 |          |   |          |   |        |
| NEMATICIDES                     |                            |         |           |           |    |          |   |          |   |        |
| Aldicarb                        | 57                         | -       | -         | 43        | 35 |          |   |          |   |        |
| Carbofuran                      | <u>c/</u>                  | -       | 70        | <u>c/</u> | 50 |          |   |          |   |        |
| Ethoprop                        | -                          | -       | <u>c/</u> | <u>c/</u> | 71 |          |   |          |   |        |
| Ethylene dibromide              | -                          | -       | -         | 39        | 39 |          |   |          |   |        |
| FUNGICIDES                      |                            |         |           |           |    |          |   |          |   |        |
| Benomyl                         | <u>c/</u>                  | 70      | -         | 70        | 45 |          |   |          |   |        |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

b/ A coefficient of variation is the standard error of the estimate divided by acres treated times 100. A coefficient of variation describes the relative variation of the estimate. The lower the value of the coefficient, the more reliable the estimate.

c/ Use of this material at the State level was not significant and was reported in the "other" or "total" category.

Appendix Table 2. Percentage of soybean herbicide acre-treatments by primary weeds controlled as reported by farmers in the Southeast, 1980 a/

|                       |       |         |    |         |       |          |   |          |   |        |
|-----------------------|-------|---------|----|---------|-------|----------|---|----------|---|--------|
|                       | :     | :       | :  | North   | :     | South    | : |          |   |        |
| Herbicides, weeds     | :     | Alabama | :  | Georgia | :     | Carolina | : | Carolina | : | Region |
|                       | ----- |         |    | Percent | ----- |          |   |          |   |        |
| <u>Alachlor</u>       |       |         |    |         |       |          |   |          |   |        |
| Grasses:              |       |         |    |         |       |          |   |          |   |        |
| Barnyardgrass         | 50    | -       | -  | -       | 9     |          |   |          |   |        |
| Broadleaf signalgrass | -     | -       | 12 | -       | 5     |          |   |          |   |        |
| Crabgrass             | 17    | 75      | 19 | 33      | 35    |          |   |          |   |        |
| Johnsongrass          | -     | -       | 13 | 17      | 8     |          |   |          |   |        |
| Panicum               | -     | -       | 6  | 33      | 7     |          |   |          |   |        |
| Other                 | -     | -       | 6  | -       | 2     |          |   |          |   |        |
| Broadleaves:          |       |         |    |         |       |          |   |          |   |        |
| Cocklebur             | 17    | 13      | 13 | -       | 12    |          |   |          |   |        |
| Morningglory          | -     | -       | 19 | -       | 8     |          |   |          |   |        |
| Pigweed               | -     | 12      | 6  | -       | 6     |          |   |          |   |        |
| Sicklepod             | 16    | -       | -  | -       | 3     |          |   |          |   |        |
| Nutsedge              | -     | -       | 6  | 17      | 5     |          |   |          |   |        |
| <u>Bentazon</u>       |       |         |    |         |       |          |   |          |   |        |
| Broadleaves:          |       |         |    |         |       |          |   |          |   |        |
| Cocklebur             | 94    | 100     | 67 | 90      | 91    |          |   |          |   |        |
| Ragweed               | 6     | -       | 17 | -       | 5     |          |   |          |   |        |
| Sicklepod             | -     | -       | -  | 10      | 2     |          |   |          |   |        |
| Other                 | -     | -       | 16 | -       | 2     |          |   |          |   |        |
| <u>Metribuzin</u>     |       |         |    |         |       |          |   |          |   |        |
| Grasses:              |       |         |    |         |       |          |   |          |   |        |
| Broadleaf signalgrass | -     | 9       | -  | -       | 4     |          |   |          |   |        |
| Crabgrass             | -     | 8       | 25 | -       | 6     |          |   |          |   |        |
| Other                 | 9     | -       | -  | -       | 3     |          |   |          |   |        |
| Broadleaves:          |       |         |    |         |       |          |   |          |   |        |
| Cocklebur             | 37    | 56      | -  | 50      | 41    |          |   |          |   |        |
| Morningglory          | 36    | -       | 25 | -       | 18    |          |   |          |   |        |
| Pigweed               | -     | -       | 25 | -       | 3     |          |   |          |   |        |
| Ragweed               | -     | -       | 25 | -       | 3     |          |   |          |   |        |
| Sicklepod             | 9     | 9       | -  | 50      | 10    |          |   |          |   |        |
| Other                 | 9     | 18      | -  | -       | 12    |          |   |          |   |        |
| <u>Trifluralin</u>    |       |         |    |         |       |          |   |          |   |        |
| Grasses:              |       |         |    |         |       |          |   |          |   |        |
| Broadleaf signalgrass | -     | -       | -  | 5       | 2     |          |   |          |   |        |
| Crabgrass             | 31    | 62      | 63 | 52      | 50    |          |   |          |   |        |
| Johnsongrass          | 37    | 8       | 13 | 3       | 15    |          |   |          |   |        |
| Panicum               | -     | -       | 12 | 3       | 2     |          |   |          |   |        |
| Other                 | 4     | 8       | -  | 2       | 5     |          |   |          |   |        |
| Broadleaves:          |       |         |    |         |       |          |   |          |   |        |
| Cocklebur             | 12    | 4       | 12 | 29      | 15    |          |   |          |   |        |
| Pigweed               | 12    | 13      | -  | 3       | 8     |          |   |          |   |        |
| Other                 | 4     | 5       | -  | 3       | 3     |          |   |          |   |        |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.



Appendix Table 3. Percentage of soybean insecticide acre-treatments by primary insects controlled as reported by farmers in the Southeast, 1980 a/

|                       |   |         |   |         |   |          |   |          |   |        |
|-----------------------|---|---------|---|---------|---|----------|---|----------|---|--------|
|                       | : | :       | : | North   | : | South    | : |          |   |        |
| Insecticides, insects | : | Alabama | : | Georgia | : | Carolina | : | Carolina | : | Region |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |
|                       |   |         |   |         |   |          |   |          |   |        |

- None reported.


a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.

Appendix Table 4. Percentage of soybean fungicide acre-treatments by primary diseases controlled as reported by farmers in the Southeast, 1980 a/

| Fungicides, diseases   | : Alabama           | : Georgia | : North Carolina | : South Carolina | : Region |
|------------------------|---------------------|-----------|------------------|------------------|----------|
|                        | ----- Percent ----- |           |                  |                  |          |
| <u>Benomyl</u>         |                     |           |                  |                  |          |
| Leaf blight            | -                   | 33        | -                | -                | 12       |
| Pod and stem blight    | 100                 | 67        | -                | 100              | 88       |
| <u>Benomyl + other</u> |                     |           |                  |                  |          |
| Anthracnose            | -                   | -         | 100              | -                | 100      |

- None reported.

a/ "1980 Soybean Pesticide Use Survey", USDA, ESS, Natural Resource Economics Division.



R0001 245132

*a*



R0001 245132